

RAYSCAN ALPHA PLUS The user can conveniently adjust the FOV according to the purpose of the treatment

02 Image Processing Technology

70µm voxel size (4cm diameter) and 6 second reconstruction time (16cm diameter)

03 Specialized for Endodontic Procedures

- Voxel size: 70µm - Minimum FOV: 4x3cm

04 One Shot Ceph

- Fast acquisition to reduce patient movement and image distortion
- Minimize patient dose with a 0.3 second exposure

05 Panoramic

The state of the art technology for high-definition image quality








The user can conveniently adjust the FOV according to the purpose of the treatment.

Dose Reduction


CBCT














01 Visible X-ray Guide

Visible X-ray Guide - Free FOV



The user gets the right image the first time and makes it more comfortable for the patient.

Consistently easy to position the patient for any image. Choose the appropriate size for any procedure. Keep the ALARA principle and reduce liability.



02 Image Processing Technology

When you need to scan faster…

Down to 4.9 second scan and 4 second 3D reconstruction!

When you need to see more detail…

Up to 70 μm with a focused field of view!

When you need to see more anatomy...

Up to 16 cm diameter field of view reconstructed in 6 seconds!









03 Specialized for Endodontic Procedures

Scan and reconstruct at 70 µm with a focused field of view for precise endodontic cases




More Detail, More Confidence, More Procedures

See more detail using a high-resolution CT image in a specific area.


Have more confidence before, during, and after the procedure with increased awareness.

Do more procedures in less time.



04 One Shot Ceph

One Shot Cephalometric Imaging acquires images in less than 1 second to reduce image distortion!



Choose from two sizes of one shot cephalometric sensors. A scanning ceph is also available for a smaller overall unit footprint.

One Shot Cephalometry

Our cutting-edge Flat Panel Detector(FPD) provides a new level of performance and reliability while reducing radiation exposure and image distortion due to patient's movement. Two different sizes of FPD are available.





Scanning Cephalometry

Our scanning ceph module allows clinicians to upgrade their diagnostic capabilities while keeping costs to a minimum.



04 One Shot Ceph

Longer exposure times can result in greater movement during acquisition of cephalometric radiographs. Because patient movement was more significant in children, a shorter exposure time is recommended in order to get a good quality cephalometric image for pediatric patients.



Small variations (1 to 2 mm) in the identification of certain landmarks can lead to different angular measurements. Regarding diagnosis, in 56% of the cases skeletal classification was changed and in 52% of the cases malocclusion classification was altered, after evaluating the LCR.




Fig 5. The amount of the subjects' movements. The youngest larger increasing proportion of the movements

Group 1: 9 to 12 years old **Group 2**: 13 to 19 years old **Group 3**: 20 to 25 years old **Group 4**: 26 to 30 years old

Group 1 represents

the main patients of orthodontic treatment and should not be imaged with a scanning ceph due to risk of patient movement.



Fig 1. The subject positions at the digital lateral cephalometric The optical marker is attached to the subject's chin.




Fig 2. Optical tracker (Polaris Vicra System, Northern Digita Waterloo, Canada).

"Quantitative Evaluation of Patient Movement during Simulated Acquisition of Cephalometric Radiographs", Kyung-Hoe Huh, Erika Benavides, Young-Tak Jo, Bo-Ram Choi, Won-Jin Yi, Min-Suk Heo, Sam-Sun Lee, and Soon-Chul Choi, Journal of Digital Imaging, Vol 24, No 3 (June), 2011: pp 552Y559

"The influence of using 2D cephalometry on orthodontic treatment outcome" Conference: 14th Congress of the European Academy of Dento-MaxilloFacial radiology, At Cluj-Napoca

05 Panoramic


The state of the art technology for high-definition image quality




FMX(Full Mouth X-ray), extracting from a panoramic image



Bitewing



Orthogonal





TMJ

Non-directional wireless remote control facilitates your patient positioning easier than ever.



07 Various options for your practice

13x10cm or 16x10cm maximum FOV options
3 different cephalometry options including One Shot Ceph

Specifications are subject to change without prior notice.

RAYSCAN α + (Model : RCT700)

Type Panoramic, Cephalometric, Cone Beam CT
Patient positioning Standing (wheelchair accessible)

Focal spot 0.5

Tube current 4~17mA

Tube voltage 60~90kVp

	α+ 160	СВСТ	Panoramic	α+ 130	СВСТ	Panoramic
Detector type		CMOS	CMOS		CMOS	CMOS
FOV / Image size		Max. 16x10cm	Max. 15cm (H)		Max. 13x10cm	Max. 14.4cm (H)
Free FOV support		Yes	Yes		Yes	Yes
Voxel size		70~400µm			70~400µm	
Exposure time		4.9~14sec	Max. 14sec		4.9~14sec	Max. 14sec

	Cephalometric (Option)					
Туре	SC (Scanning Ceph)	OCL (One shot Large)	OCS (One shot Standard)			
Detector type	CdTe detector	a-Si TFT	a-Si TFT			
Image size	Max. 26x24cm	Max. 33x33cm	Max. 30x25cm			
Exposure time	4.8~10.4sec	0.3 / 0.8sec	0.3 / 0.8sec			

IDEA BRONZE

INTERNATIONAL DESIGN EXCELLENCE AWARDS 112

REDDOT WINNER

reddot design award winner 2012

GD BEST OF BEST





RBS-AP03 (rev.1)

Design and specifications are subject to change without notice

Ray Co., Ltd.

332-7, Samsung1-ro, Hwaseong-si, Gyeonggi-do, 18380, Korea

Phone +82.31.605.1000

Email ray_overseas@raymedical.co.kr

Web www.raymedical.com Ray Digital solution

For more information or inquiries, please visit www.raymedical.com or contact your local representative.







